



## High Energy Creep 18+ R40

### MEDICATED

For Growing cattle on pasture or in dry lot

#### Active Drug Ingredient

Monensin..... 40g/ton (20 mg/lb)

#### Guaranteed Analysis

Crude Protein (Min).....	18%
Crude Fat (Min).....	4.75%
Fiber (Max).....	20%
Calcium (Min).....	.90% <sup>b</sup>
Calcium (Max).....	1.25%
Phosphorus (Min).....	.50%
Salt (Min) .....	.40%
Salt (Max) .....	.70%
Potassium (Min).....	.75%
Vitamin A (Min).....	5,700 IU/LB

#### Ingredient Statement

Processed Grain By-Products, Plant Protein Products, Roughage Products, Calcium Carbonate, Salt, Yeast Culture, Propionic Acid (as preservative), Vitamin A Supplement, Vitamin E Supplement, Ferrous Sulfate, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Copper Chloride, Ethylenediamine Dihydriodide, Cobalt Carbonate, Copper Amino Acid Complex, Zinc Amino Acid complex, Vitamin D3 Supplement, and Calcium Iodate

#### Feeding Directions

For the first 5 days, feed at the rate of 2.5 to 5 lbs per head per day to provide 50 to 100 mg of Monensin. After 5 days can feed up to 10 lbs per head per day. Not to exceed 10 lbs per head per day. Do not self-feed. A roughage source or pasture should be available. Provide plenty of fresh, clean water at all times.

**CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are red to other groups of cattle, the concentration of Monensin in the refusals are amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to**

**be processed for veal.**

Henry's Healthy Harvest  
246 Basher Drive, Unit 7  
Berthoud, CO 80513  
(970)308-7332

