



Performance Goat Feed

For prevention of coccidiosis in goats caused by Eimeria Crandalis, E. Christenseni and E. Ninakohlyakimovae

Active Drug Ingredients

MONENSIN..... 20g/ton(10mg/lb)

Guaranteed Analysis

Crude Protein (Min).....	18.0%*
*Includes not more than 1.2% equivalent from non-protein nitrogen	
Crude Fat (Min).....	7.0%
Crude Fiber (Max).....	19.0%
Calcium (Min).....	.90%
Calcium (Max).....	1.25%
Phosphorus (Min).....	.40%
Salt (Min)20%
Salt (Max)	0.70%
Copper (Min)	30 ppm
Copper (Max)	60 ppm
Selenium (Min)30 ppm
Vitamin A (Min)	4800 IU/LB

Ingredient Statement

Processed Grain By-Products, Plant Protein Products, Grain Products, Roughage products, Calcium Carbonate, Vegetable Fat, Molasses Products, Ammonium Chloride, Salt, Propionic acid(as preservative), Yeast Culture, Vitamin D3 Supplement, Ferrous Sulfate, Zinc Sulfate, Copper Amino Acid Complex, Zinc Amino Acid Complex, Copper Sulfate, Calcium Lodate, Sodium Selenite, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Copper Chloride, Ethylenediamine Dihydriodide, and Cobalt Carbonate.

Feeding Directions

Feed continuously as the sole ration to goats in confinement at the rate of 2-3% of the growing goats body weight. Offer quality forage. Provide plenty of fresh water at all times.

DO NOT feed to sheep or wool producing goats

CAUTION: For Ruminants (goats) only. Do not allow horses access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

Henry's Healthy Harvest
246 Basher Drive, Unit 7
Berthoud, CO 80513
(970)308-7332